

## Meningococcal B Immunisation Program

Administration of paracetamol with every dose of Bexsero® is recommended for children aged less than 2 years. Fever is a common reaction following any vaccine but it is more common with Bexsero. Paracetamol will reduce the likelihood of high-grade fever for your child.

Parents are recommended to give the **1st dose of paracetamol 30 minutes before immunisation**, or as soon as possible after immunisation. Give 2 more doses of paracetamol 6 hours apart, even if your child does not have a fever.

Dose 1 (30 minutes before, or as soon as possible after immunisation)	Time given ____:____
Dose 2 (6 hours from 1st dose)	Time given ____:____
Dose 3 (6 hours from 2nd dose)	Time given ____:____

## Immunisation record (to be completed by doctor or nurse)

My name \_\_\_\_\_ Date of birth \_\_\_\_ / \_\_\_\_ / \_\_\_\_ Medicare number \_\_\_\_\_

Age	Immunisation	Batch number	Date given	Next due	Provider signature / stamp
6 weeks	Meningococcal B LL				
4 months	Meningococcal B LL				
12 months	Meningococcal B LA				

If your child does develop a fever and you feel the fever is not adequately controlled using the above recommendations you should refer to the dosing schedule on the paracetamol bottle or seek further medical advice.